

Logarithmic Discriminants of Hyperplane Arrangements

Regiomontanus PhD Seminar


Leonie Kayser, Andreas Kretschmer & Simon Telen

leokayser.github.io

October 23, 2024



MAX PLANCK INSTITUTE
FOR MATHEMATICS
IN THE SCIENCES



Duolingo Algebraic Geometry

- ▷ $g_1, \dots, g_s \in \mathbb{C}[x_0, \dots, x_n]$ polynomials, then

$$X = V(g_1, \dots, g_s) := \{ x \in \mathbb{C}^{n+1} \mid g_1(x) = \dots = g_s(x) = 0 \}$$


- ▷ X is called an *algebraic variety*, a (possibly singular) complex submanifold of \mathbb{C}^{n+1}
- ▷ X is irreducible if it is not a proper union of varieties
- ▷ $\mathbb{P}^n = (\mathbb{C}^{n+1} \setminus 0) / \sim$, $x \sim y$ iff $x = \lambda y$ for some $\lambda \in \mathbb{C}^\times := \mathbb{C} \setminus 0$, home of homogeneous polynomials
- ▷ Hypersurface = variety defined by a single equation h , degree is the degree of X
- ▷ X smooth if $J_g(x)$ has maximal rank for all $x \in X$
- ▷ x is a non-reduced/degenerate solution to g_1, \dots, g_s if the Jacobi matrix $J_g(x)$ does not have maximal rank

Maximum Likelihood Estimation in Algebraic Statistics

- ▷ Let $X \subseteq (\mathbb{C}^\times)^{n+1}$ be a d -dimensional smooth very affine variety
- ▷ Discrete statistical model $X \cap \Delta_n = \{ p \in X \cap \mathbb{R}^{n+1} \mid p_i > 0, p_0 + \dots + p_n = 1 \}$
- ▷ Given data points $u \in \mathbb{N}^{n+1}$, which parameter maximizes the log-likelihood

$$\mathcal{L}_u(x) = \log x_0^{u_0} \cdots x_n^{u_n}, \quad x \in X \cap \Delta_n?$$

- ▷ Critical equations: $x \in \text{Crit}_X(u) = \{ x \in X \mid \nabla \mathcal{L}_u(x) = 0 \}$
- ▷ $\text{Crit}_X(u)$ is a finite set of $\text{MLdeg}(X)$ non-degenerate critical points for general data $u \in \mathbb{N}^{n+1}$ (or \mathbb{C}^{n+1})
- ▷ [Huh13] $\text{MLdeg}(X) = (-1)^d \cdot \chi(X)$
- ▷ Extensively studied for toric models (exponential families), linear models, determinantal varieties, ...



Linear models and scattering amplitudes

- ▶ Let be an essential arrangement of $n + 1$ hyperplanes in \mathbb{C}^d

$$\mathcal{A} = \mathbb{V}(\ell_0 \cdots \ell_n) \subseteq \mathbb{C}^d, \quad (\ell_0(x), \dots, \ell_n(x))^T = Ax + b, \quad L^T = [b \mid A]$$

- ▶ Parametrizes linear model $X := \mathbb{C}^d \setminus \mathcal{A} \xrightarrow{\ell} (\mathbb{C}^\times)^{n+1}$, can assume $\sum_j \ell_j = 1$
- ▶ Log-likelihood function or master function given by

$$\mathcal{L}_u(x) = u_0 \log \ell_0(x) + \cdots + u_n \log \ell_n(x), \quad \nabla \mathcal{L}_u(x) = A^T \text{diag}(1/\ell_0, \dots, 1/\ell_n)u$$

- ▶ If the ℓ_j are real, then $\text{MLdeg}(X)$ is the number of bounded chambers of $\mathcal{A} \cap \mathbb{R}^d$
- ▶ Critical equations appear as scattering equations in bi-adjoint scalar ϕ^3 -theories (Cachazo, He & Yuan [CHY14])

What is “general” data?

- ▷ Moving from general to special $u \in \mathbb{P}^n = \mathbb{P}(\mathbb{C}^{n+1})$, what can happen to $\text{Crit}_X(u)$?
 1. Two critical points collide to form a non-reduced/degenerate point
 2. A positive-dimensional component appears
 3. A critical point disappears to infinity
- ▷ The closure of 1.-3. was called the *data discriminant* in [RT15]
- ▷ 3. was studied by (Sattelberger & van der Veer [SvdV23])

Definition

The *logarithmic discriminant* of a (smooth) variety $X \hookrightarrow (\mathbb{C}^\times)^{n+1}$ is

$$\nabla_{\log}(X) := \overline{\{u \in \mathbb{P}^n \mid \text{Crit}_X(u) \text{ is infinite or non-reduced}\}}.$$

↪ Goal: Understand logarithmic discriminants of hyperplane arrangements!

Three points enter a bar

- ▶ Three points on a line $\mathcal{A} = V(x(x+1)(x+b)) = \{0, -1, -b\} \subseteq \mathbb{C}^1$ ($b \notin \{0, 1\}$)
- ▶ Model is a line $X \subseteq (\mathbb{C}^\times)^3$ parametrized by $(x, x+1, x+b)$,

$$\mathcal{L}_u(x) = u_0 \log x + u_1 \log(x+1) + u_2 \log(x+b)$$

- ▶ A single critical equation in $x \in \mathbb{C}^1 \setminus \mathcal{A}$

$$\frac{u_0}{x} + \frac{u_1}{x+1} + \frac{u_2}{x+b} = 0 \iff u_0(x+1)(x+b) + u_1x(x+b) + u_2x(x+1) = 0$$

- ▶ When does this quadric have a double root in x ? Highschool discriminant!

$$\Delta_{\log}(X) = (b-1)^2 u_0^2 + 2b(b-1) u_0 u_1 + b^2 u_1^2 - 2(b-1) u_0 u_2 + 2b u_1 u_2 + u_2^2$$

- ▶ $\Delta_{\log}(X)$ itself is a smooth quadric in u with discriminant $-4b^2(b-1)^2$

Ramification and its consequences

- ▷ Let $f: V \rightarrow W$ be a dominant map of smooth irreducible varieties of dimension n
- ▷ The ramification locus $\text{Ram}(f) \subseteq V$ is the hypersurface

$$\text{Ram}(f) = \{ x \in V \mid x \in f^{-1}(f(x)) \text{ is not isolated or reduced} \} = V(\det J_f(x))$$

- ▷ The branch locus is the image closure $\text{Branch}(f) = \overline{f(\text{Ram}(f))} \subseteq W$
- ▷ Apply this to the *likelihood correspondence*

$$f: \mathcal{L}_X^\circ = \{ (u, x) \in \mathbb{P}^n \times X \mid \nabla \mathcal{L}_u(x) = 0 \} \rightarrow \mathbb{P}^n$$

Definition (True definition of $\nabla_{\log}(X)$)

The logarithmic discriminant is the branch locus of the projection f . The ramification locus is defined in $\mathbb{P}^n \times X$ by

$$\nabla \mathcal{L}_u(x) = 0, \quad \det \text{Hess}_x(\mathcal{L}_u(x)) = 0.$$

Irreducibility of $\text{Ram}(f)$

- ▷ $X = \mathbb{C}^d \setminus V(\ell_0 \cdots \ell_n)$, $(\ell_0(x), \dots, \ell_n(x))^T = Ax + b$
- ▷ Here the equations of the ramification locus have a very concrete form

$$\nabla \mathcal{L}_u(x) = A^T \cdot \text{diag}(1/\ell_0, \dots, 1/\ell_n) \cdot u = 0$$

$$h = \det \left(A^T \cdot \text{diag} \left(\frac{u_0}{\ell_0^2}, \dots, \frac{u_n}{\ell_n^2} \right) \cdot A \right) = \sum_{\substack{I \subseteq \{0, \dots, n\} \\ |I|=d}} |A_I|^2 \frac{u^I}{(\ell^I)^2}$$

- ▷ Critical equations are linear in the u_j \rightsquigarrow substitute them in h to obtain

$$\tilde{h} \in \mathbb{C}[u_d, \dots, u_n; x], \quad \text{Ram}(f) \cong V(\tilde{h}) \subseteq \mathbb{P}^{n-d} \times X$$


Theorem

If the arrangement contains a subset of $d + 2$ hyperplanes which is bi-uniform (to be defined in a moment), then $\text{Ram}(f)$ and hence $\nabla_{\log}(X)$ are irreducible varieties.

A split discriminant!

- ▶ Consider the arrangement \mathcal{A} of six planes

$$(1, x_1, x_2, x_3) \cdot \begin{bmatrix} 1 & 2 & 1 & 0 & 0 & 0 \\ \hline 1 & 1 & 2 & 1 & 0 & 1 \\ 1 & \frac{3}{2} & \frac{3}{2} & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 & 1 & 2 \end{bmatrix}$$



- ▶ The first and the last three planes intersect in a line each
- ▶ The logarithmic discriminant decomposes as

$$\begin{aligned} \nabla_{\log} = & V(144u_0^2 + 120u_0u_1 + 168u_0u_2 + 25u_1^2 - 70u_1u_2 + 49u_2^2) \\ & \cup V(u_3^2 - 2u_3u_4 + 4u_3u_5 + u_4^2 + 4u_4u_5 + 4u_5^2) \\ & \cup V(u_0 + u_1 + u_2, u_3 + u_4 + u_5). \end{aligned}$$

Theorem

Let $\mathcal{A} \subseteq \mathbb{C}^1$ be an arrangement of $n + 1 \geq 3$ distinct points.

1. The ramification locus is a smooth irreducible hypersurface in $\mathbb{P}^n \times (\mathbb{C}^1 \setminus \mathcal{A})$.
2. The class of $\overline{\text{Ram}(f)}$ in the cohomology ring $H_{\text{sing}}^{2\bullet}(\mathbb{P}^n \times \mathbb{P}^1) = \mathbb{Z}[\alpha, \beta]/\langle \alpha^{n+1}, \beta^2 \rangle$ is $\alpha^2 + 2(n - 1)\alpha\beta$.
3. The projection $f: \text{Ram}(f) \rightarrow \nabla_{\log}$ is generically bijective.
4. $\nabla_{\log}(X) \subseteq \mathbb{P}^n$ is an irreducible hypersurface of degree $2(n - 1)$.

▷ Explicit formula

$$\Delta_{\log} = \text{Disc}_x \left(\sum_{i=0}^n u_i \prod_{k \neq i} (x + b_k) \right)$$

▷ For $n + 1 = 4$ points $\nabla_{\log} \subseteq \mathbb{P}^3$ is always a singular surface of degree 4

General arrangements

- ▷ \mathcal{A} defined by $(\ell_0(x), \dots, \ell_n(x))^T = Ax + b$
- ▷ $k \times n$ matrix is *uniform* if all sets of k columns are linearly independent
- ▷ *Little matroid* $M(A^T)$, *big matroid* $M([b|A]^T)$, bi-uniform = both are uniform
- ▷ $\nabla_{\text{Hu}}(\mathcal{A}) \supseteq \nabla_{\text{log}}(\mathcal{A})$ is a second kind of discriminant

Theorem

Let \mathcal{A} be a bi-uniform arrangement of $n + 1 \geq d + 2$ hyperplanes in \mathbb{C}^d .

1. $\nabla_{\text{Hu}}(\mathcal{A})$ is a hypersurface of degree $2d \binom{n-1}{d}$ with full Newton polytope
2. $\nabla_{\text{log}}(\mathcal{A})$ is an irreducible and reduced hypersurface.
3. $\nabla_{\text{log}} \subseteq \nabla_{\text{Hu}}$ coincide as sets, so $\nabla_{\text{Hu}} = V((\Delta_{\text{log}})^e)$ for some $e \geq 1$.
4. If the arrangement is defined by real affine linear forms, then $\nabla_{\text{log}} \cap \mathbb{R}_+^{n+1} = \emptyset$.

- ▷ We know that equality hold for $d = 1$ and expect this to always hold true

Many open questions

- ▷ Missing piece of the puzzle: Is ∇_{Hu} reduced for a bi-uniform arrangement?
- ▷ (When) is the projection $\text{Ram}(f) \rightarrow \nabla_{\log}$ generically one-to-one? ... bijective?
- ▷ Is there any arrangement such that $\nabla_{\log}(\mathcal{A})$ is *not* reduced?
- ▷ Is there an arrangement of lines whose logarithmic discriminant is reducible?
- ▷ Is the degree of $\nabla_{\log}(\mathcal{A})$ an invariant of the little and big matroid?
- ▷ What is the meaning of the components $\nabla_{\text{Hu}} \setminus \nabla_{\log}$?

Thank you! Questions?

arXiv:2410.11675

The discriminant of $\mathcal{M}_{0,m}$

- ▷ $\mathcal{M}_{0,m}$ parametrizes tuples of m points on the projective line \mathbb{P}^1
- ▷ By fixing $(0, 1, x_1, \dots, x_{m-3}, \infty)$, it can be realized as the complement in \mathbb{C}^{m-3} of the $n = \binom{m-1}{2} - 1$ minors of

$$\begin{pmatrix} 1 & 1 & 1 & 1 & \cdots & 1 & 0 \\ 0 & 1 & x_1 & x_2 & \cdots & x_{m-3} & 1 \end{pmatrix}$$




- ▷ Variable corresponding to minor (i, j) are *Mandelstam invariants* s_{ij}
- ▷ Discriminant for $m = 5$ has degree $4 < 2 \cdot 2 \cdot \binom{5-2}{2} = 12$

$$\Delta_{\log}(\mathcal{M}_{0,5}) = (s_{13}s_{24} + s_{13}s_{34} + s_{14}s_{34} + s_{14}s_{23} + s_{23}s_{34} + s_{24}s_{34} + s_{34}^2)^2 - 4s_{13}s_{14}s_{23}s_{24}$$

- ▷ The Hurwitz discriminant has the extra factors

$$\Delta_{\text{Hu}}(\mathcal{M}_{0,5}) = (s_{13} + s_{23} + s_{34})^2 \cdot (s_{14} + s_{24} + s_{34})^2 \cdot \Delta_{\log}(\mathcal{M}_{0,5})$$

- ▷ Conjecturally rich nested structure, degrees of $\nabla_{\log}(\mathcal{M}_{0,m})$ are $4, 30, 208, 1540, \dots$

-  Freddy Cachazo, Song He, and Ellis Ye Yuan.
Scattering equations and Kawai-Lewellen-Tye orthogonality.
Physical Review D, 90(6):065001, 2014.
-  June Huh.
The maximum likelihood degree of a very affine variety.
Compositio Mathematica, 149(8):1245–1266, 2013.
-  Leonie Kayser, Andreas Kretschmer, and Simon Telen.
Logarithmic discriminants of hyperplane arrangements, 2024.



Jose Israel Rodriguez and Xiaoxian Tang.

Data-discriminants of likelihood equations.

In *Proceedings of the 2015 ACM on international symposium on symbolic and algebraic computation*, pages 307–314, 2015.



Anna-Laura Sattelberger and Robin van der Veer.

Maximum likelihood estimation from a tropical and a Bernstein–Sato perspective.

International Mathematics Research Notices, 2023(6):5263–5292, 2023.